

## SHORTACTIVE DRY

### DRY ACIDIFIER WITH BUTYRIC ACID

#### FEED FOR ALL SPECIES

Contains: calcium carbonate, monoglycerides, diglycerides and triglycerides of butyric acid, glycerol.

ADDITIVES (per kg)

- Preservatives:

E260 Acetic acid mg 35,800 - E270 Lactic acid mg 10,200 - E238 Calcium formate mg 150,000

E282 Calcium propionate mg 130,000 - E338 Orthophosphoric acid mg 18,600

Binders: E551a precipitated and dried silicic acid mg 140,000

Anti-caking agents: 1g568 Clinoptilolite of sedimentary origin mg 207.600

#### DESCRIPTION

The organic acids present in the product prevent the formation of mold and the development of toxic bacteria in the gastrointestinal tract of animals.

It helps to keep the pH of the gastrointestinal tract at optimal values.

Thanks to the presence of protected butyric acid monodiglycerides and the effects of all its ingredients, SHORTACTIVE DRY improves the health of the digestive system and intestines in terms of:

- Increased intake and reduction of bacterial content of feed
- Improvement of microflora and development of a symbiotic microbiota
- Control of the development of pathogenic bacteria
- Reduction of inflammatory cells

#### ACTION

SHORTACTIVE DRY is an acidifier that stabilizes the feed and controls the development of mold.

In addition, the mixture of p.a. acidifiers, combined with Butyric Acid (protected), allows to exploit a natural synergism between the components, developing a strong action in the intestine.

In adult animals it exerts bacteriostatic effect.

#### DOSAGE

To be mixed thoroughly in complete feed at the dose of:

poultry: 1 - 2.5 kg / ton of complete feed.

Store in a cool and dry place, away from heat sources.

Dosages may vary depending on the company's health situation.

The technical service is available to recommend the dosage.

#### PACKAGES

Available in bags of kg.25 - Big Bag of kg.1000

Store in a cool, dry place, away from heat sources.

Consume within 12 months from the date of manufacture.

FAMI-QS Certificate